

Benefit[®] 60 WP

**Greenhouse and Nursery Insecticide in
Water Soluble Packaging**
For Foliar and Systemic Insect Control on Ornamental
and Vegetable Plants in Greenhouses, Nurseries and
Interior Plantscapes

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl) methyl]-
N-nitro-2-imidazolidinimine 60%
OTHER INGREDIENTS: 40%

Total 100%

EPA Reg. No. 42750-153-58185 EPA Est. 65387-AR-002

Keep water soluble packets in this container and store in a cool dry place but not below freezing (32°F).

Do Not Remove Packets From Container Except For Immediate Use.

**STOP - READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

In case of emergency call toll free at 1-888-241-6409. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

This specimen label is for information only.
Everris makes no representations as to the accuracy of this label. It is the responsibility of the user to read and follow the label attached to the pesticide product container.

**NET CONTENTS: 3.53 oz. (100 g)
[Five - 0.71 oz. (20 g) packets]**

Stock # E956601

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on the foliage of blooming crops, plants or weeds. Do not apply this product or allow it to drift to blooming crops, plants or weeds if bees are foraging the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS exist for this product because of risk to bees and other insect pollinators. Follow application restrictions found in the directions for use to protect pollinators.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not formulate this product into other end-use products.

See individual crops for specific pollinator protection application restrictions. If none exist under the specific crop, for foliar applications, follow these application directions for crops that are contracted to have pollinator services or for food/feed & commercially grown ornamentals that are attractive to pollinators.



1. FOR FOOD CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.



2. NON-AG USE SITES:

Do not apply Benefit 60 WP while bees are foraging. Do not apply Benefit 60 WP to plants that are flowering. Only apply after all flower petals have fallen off.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep children and pets off treated area until dry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

USE INFORMATION

BENEFIT® 60 WP is for insect control on ornamental and vegetable plants in nurseries, greenhouses and interior plantscapes. BENEFIT 60 WP is a systemic product and will be translocated upward within the plant. To assure optimum effectiveness, the product must be placed where the roots of the plant can absorb the active ingredient.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Irrigate moderately but thoroughly after application, allowing no leaching and runoff from container for at least three irrigations or 10 days whichever is longer.

Resistance: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area. For resistance management purposes, a foliar application of any chloronicotinyl insecticide following a BENEFIT 60 WP soil application in the same crop is not recommended.

Incorporation: Incorporation of BENEFIT 60 WP can be achieved by cultivation, irrigation, rainfall, mechanical placement, soil injection, drenching, and broadcast sprays.

Woody Perennials: Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks should be expected. Longer delays may be expected with larger plants. Application should therefore be made well in advance of expected insect activity.

Bark Media: Media with 30 to 50% or more bark content may confer a shorter period of protection when treated with BENEFIT® 60 WP.

Tank Mixes: BENEFIT 60 WP has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

Do not use BENEFIT 60 WP packets in a tank mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water. However, normal chlorinated water is acceptable for mixing.

MIXING: Within each foil pouch is a clear inner packet containing BENEFIT 60 WP. The clear inner packet is water soluble. Do not allow packets to become wet prior to adding to the tank. Do not handle the clear inner packets with wet hands or wet gloves. Rough handling may cause breakage.

Reseal outer carton to protect remaining packets. To prepare the mixture, remove the outer foil pouch and drop the required number of unopened clear water soluble packets, as determined under "Recommended Drench and Irrigation Applications", into the spray tank while filling with water to the desired level. Operate the agitator while mixing.

Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within a few minutes from the time they are added to the water. Cooler water temperatures increase the time needed for the inner packet to dissolve completely.

APPLICATION THROUGH IRRIGATION SYSTEMS

BENEFIT 60 WP may be applied at rates recommended on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:10 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply BENEFIT® 60 WP only through micro-irrigation (individual spaghetti tube), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

APPLICATION TO GRASSY AREAS AROUND NURSERIES

BENEFIT 60 WP can be used for the control of the following soil inhabiting pests of grassy areas of nurseries:

Northern and Southern masked chafers,
Cylocephala borealis, *C. immaculata*, and/or *C. lurida*;
 Asiatic garden beetle, *Maladera castanea*;
 European chafer, *Rhizotroqus majalis*;
 Green June beetle, *Cotinis nitida*;
 May or June beetle, *Phyllophaga spp.*;
 Japanese beetle, *Popillia japonica*;
 Oriental beetle, *Anomala orientalis*;
 Billbugs, *Spherophorus spp.*;
 Annual bluegrass weevil, *Hyperodes spp.*;
 Black turfgrass *ataenius*, *Ataenius spretulus* and *Aphodius spp.*
 Mole crickets, *Scapteriscus spp.*

BENEFIT 60 WP can also be used for suppression of cutworms and hairy chinchbugs.

BENEFIT 60 WP can be used as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

The active ingredient in BENEFIT 60 WP has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

Application Equipment for Use on Grassy Areas in Nurseries:

Apply BENEFIT 60 WP in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

RESTRICTIONS FOR GRASSY AREA:

- Application cannot exceed a total of 10.7 oz (0.4 lb of active ingredient) per acre per year.
- BENEFIT 60 WP cannot be used on commercial sod farms.
- Do not apply when grassy areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- Do not apply this product in the state of Oregon, by any application method, to linden, basswood or other *Tilia* species.

APPLICATIONS TO GRASSY AREAS OF FIELD & FOREST NURSERIES

PEST	DOSAGE
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms (suppression) European Chafer Green June beetle Japanese beetle Northern masked chafer Oriental beetle Phyllophaga spp. Southern masked chafer	One packet per 2,900-3,850 sq. ft.
Chinch bugs (suppression) Mole Crickets	One packet per 2,900 sq. ft.
<p>For optimum control of grubs, billbugs and annual bluegrass-weevil, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.</p> <p>For suppression of chinch bugs, make application prior to the hatching of the first instar nymphs. For control of mole crickets, make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, BENEFIT® 60 WP application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.</p>	
<p>Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.</p> <p><i>For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch.</i></p> <p>RESTRICTIONS FOR FIELD & FORESTRY NURSERIES</p> <ul style="list-style-type: none"> • Do not apply more than 10.7 oz. (0.4 lb of active ingredient) per acre per year. • Do not mow grass area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected. 	

APPLICATIONS FOR USE ON ORNAMENTALS

For foliar and systemic insect control in and around field-grown nursery and container stock, indoor and outdoor ornamentals (including both greenhouse and interior plantscapes) and ornamentals grown in flats, benches or beds.

SITE	PEST	DOSAGE
Trees (including non-bearing fruit and nut) Shrubs Evergreens Flowers Foliage plants Ground covers Vegetable plants*	Adelgids Aphids Japanese beetles (adults) Lacebugs Leafbeetles (including Elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies	One packet per 100 gal water
<p>Foliar Applications: Start treatments prior to establishment of high pest populations and reapply on an as needed basis. For resistance management purposes, a BENEFIT 60 WP foliar application following a soil application in the same crop is not recommended.</p>		
	White grub larvae (such as Japanese beetle larvae, chafers, Phyllophaga spp., Asiatic garden beetle, Oriental beetle)	One packet per 2,900-3,850 sq. ft.
<p>Broadcast Applications: Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1,000 sq. ft. For optimum control, irrigate thoroughly to incorporate BENEFIT 60 WP into the upper soil profile.</p> <p>Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.</p> <p>*Only for use on vegetable plants intended for resale including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.</p>		

Do not apply this product in the state of Oregon, by any application method, to linden, basswood or other *Tilia* species

DRENCH AND IRRIGATION RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not apply to soils which are water-logged or saturated, which will not allow the penetration of the insecticide into the root zone of the plants.
- Do not allow leachate runoff for the first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient.
- For outdoor ornamentals grown in beds or turf, applications of BENEFIT® 60 WP cannot exceed a total of 10.7 oz. (0.4 lb of active ingredient) per acre per year.
- On plants with a production cycle of less than one year, application is not to exceed a frequency of more than once each 16 weeks for a particular plant. On stock plants and woody crops with a production cycle of greater than one year, application may not exceed once a year.
- Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.
- For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval should be observed.
- Do not apply this product in the State of Oregon by any application method, to linden, basswood, or other *Tilia* species.

SOIL INJECTION AND DRENCH APPLICATIONS FOR NURSERY, GREENHOUSE AND INTERIORESCAPE PLANTS

Adelgids
Aphids
Armored scales (suppression)
Black vine weevil larvae
Eucalyptus longhorned borers
Flatheaded borers (including bronze birch and alder borers)
Japanese beetles (adults)
Lacebugs
Leaf beetles (including elm and viburnum leaf beetles)
Leafhoppers (including glassy-winged sharpshooter)
Leafminers
Mealybugs
Pine Tip moth larvae
Psyllids
Royal palm bugs
Sawfly larvae
Soft scales
Thrips (suppression)
White grub larvae
Whiteflies

TREES: 1 packet (0.71 oz. or 20 g) per 8 to 16 inches of cumulative trunk diameter.

Soil Injection:

- **GRID SYSTEM:** Holes should be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree.
- **CIRCLE SYSTEM:** Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.
- **BASAL SYSTEM:** Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.

Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.

No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.

Do not apply this product in Oregon, by any application method, to linden, basswood or other *Tilia* species.

Soil Drench:

Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

For Control of Specified Borers:

Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.

SHRUBS: 1 packet (0.71 oz. or 20 g) per 8 to 16 feet of cumulative shrub height.

Soil Injection:

Apply to individual plants using dosage indicated.

Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.

No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.

Soil Drench:

Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the shrub, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

FLOWERS AND GROUND COVERS: 1 packet (0.71 oz. or 20 g) per 3,000 sq. ft.

Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established prior to bloom or after petal fall is complete. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.

Restrictions for Soil Injection and Drench Application for Nursery, Greenhouse and Flowers and Ground Covers

- Do not apply more than 15 packets (300 grams) (0.4 lbs AI) per acre per year
- Do not harvest or consume fruits or nuts from trees that have been treated within 1 year of application
- Do not apply this product in the state of Oregon, by any application method, to linden, basswood or other *Tilia* species.

EBB & FLOOD APPLICATION TO ORNAMENTAL AND VEGETABLE PLANTS

BENEFIT® 60 WP may be applied through Ebb and Flood application to Ornamental and Vegetable Plants (intended for resale only) grown in containers. To assure accurate, that prior to treatment, bring a minimum of 10 plants up to a known field capacity and allow to dry out for one to two days. Re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

Adelgids
 Aphids
 Armored scales (suppression)
 Fungus Gnats (larvae only)¹
 Japanese Beetles (adults)
 Lacebugs
 Leaf beetles (including elm and viburnum leaf beetles)
 Leafhoppers (including glassy-winged sharpshooter)
 Leafminers
 Mealybugs
 Psyllids
 Root mealybugs²
 Root Weevil Complex: (such as Apopka Weevil, Black Vine Weevil, Citrus Root Weevil)³
 Soft Scales
 Thrips (suppression)⁴
 Whiteflies
 White Grub larvae (such as Japanese Beetle, Masked Chafer, European Chafer, Oriental Beetle, Asiatic Garden Beetle)

Pot sizes (inches)	Number of pots treated with one packet	
	Herbaceous species including vegetable plants ⁵ (one or two plants per pot)	Woody perennials, Herbaceous species including vegetable plants ⁵ (3 or more plants per pot)
2	3000	2000
3	2000	1350
4	1500	1000
5	1200	800
6	1000	650
7	850	550
8	750	500
9	675	450
10	600	400
11	550	350
12	500	300

Footnotes:

- ¹ Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of BENEFIT 60 WP from a healthy root system translocating the active ingredient up into the plant.
- ² Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1 packet (0.71 oz. or 20 g) in 150 gallons of water.
- ³ Citrus Root Weevil: For use on non-bearing citrus nursery stock.
- ⁴ Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.
- ⁵ For use on the following vegetable plants intended for resale only: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

Ebb & Flood Restrictions:

-Use only on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Callards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo and Tomatoes.

DISCLAIMER AND LIMITATION OF LIABILITY IMPORTANT NOTICE FROM EVERRIS NA INC. ("Everris"). PLEASE READ BEFORE USE.

To the extent consistent with applicable law, user or buyer accepts the conditions, disclaimer of warranties and limitations of liability. Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once for full refund.

CONDITIONS: The directions for use of this product are believed to be adequate and the user or buyer must always follow the label directions carefully and exercise judgment and caution when using this product under their growing conditions. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness, unsatisfactory or substandard results or other unintended consequences may result because of such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Everris. All such risks shall be assumed by the user or buyer.

WARRANTY: This product corresponds to all claims and descriptions set forth on the label and is reasonably fit for the purposes set forth in the directions for use on the label when used in accordance with those directions. This warranty is subject to the conditions and limitations stated herein. Everris recognizes that the rights and remedies of the user or buyer are subject to the provisions of the applicable law, but makes no other warranties or representations, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Everris is authorized to make any warranties beyond those contained herein or to modify the warranties contained therein. Subject to the user's or buyer's rights and remedies under the applicable law, Everris disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: Subject to the user's or buyer's rights and remedies under the applicable law, the exclusive remedy of the user or buyer and the liability of Everris or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid by the user or buyer for the quantity of this product involved or, at Everris' election, the replacement of the product. To the extent consistent with applicable law, Everris must have prompt notice of any claim so that a timely investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify Everris of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

Distributed by:
 Everris NA Inc.
 PO Box 3310
 Dublin, OH 43016
 1-800-492-8255